

## Tilburg University

### BGGM

Williams, Donald R.; Mulder, Joris

*Published in:*  
The Journal of Open Source Software

*DOI:*  
[10.21105/joss.02111](https://doi.org/10.21105/joss.02111)

*Publication date:*  
2020

*Document Version*  
Publisher's PDF, also known as Version of record

[Link to publication in Tilburg University Research Portal](#)

*Citation for published version (APA):*  
Williams, D. R., & Mulder, J. (2020). BGGM: Bayesian Gaussian Graphical Models in R. *The Journal of Open Source Software*, 5(51), [2111]. <https://doi.org/10.21105/joss.02111>

#### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

#### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

# BGGM: Bayesian Gaussian Graphical Models in R

Donald R. Williams<sup>1</sup> and Joris Mulder<sup>2</sup>

<sup>1</sup> Department of Psychology, University of California, Davis <sup>2</sup> Department of Methodology and Statistics, Tilburg University

DOI: [10.21105/joss.02111](https://doi.org/10.21105/joss.02111)

## Software

- [Review](#) ↗
- [Repository](#) ↗
- [Archive](#) ↗

Editor: [Anisha Keshavan](#) ↗

## Reviewers:

- [@jayrobwilliams](#)
- [@paulgovan](#)

Submitted: 15 January 2020

Published: 21 July 2020

## License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#)).

## BGGM: Bayesian Gaussian Graphical Models

The R package **BGGM** provides tools for making Bayesian inference in Gaussian graphical models (GGM). The methods are organized around two general approaches for Bayesian inference: (1) estimation and (2) hypothesis testing. The key distinction is that the former focuses on either the posterior or posterior predictive distribution (Gelman, Meng, & Stern, 1996; see section 5 in Rubin, 1984), whereas the latter focuses on model comparison with the Bayes factor (Jeffreys, 1961; Kass & Raftery, 1995).

## What is a Gaussian Graphical Model ?

A Gaussian graphical model captures conditional (in)dependencies among a set of variables. These are pairwise relations (partial correlations) controlling for the effects of all other variables in the model.

## Applications

The Gaussian graphical model is used across the sciences, including (but not limited to) economics (Millington & Niranjana, 2020), climate science (Zerenner, Friederichs, Lehnertz, & Hense, 2014), genetics (Chu, Weiss, Carey, & Raby, 2009), and psychology (Rodriguez, Williams, Rast, & Mulder, 2020).

## Overview

The methods in **BGGM** build upon existing algorithms that are well-known in the literature. The central contribution of **BGGM** is to extend those approaches:

1. Bayesian estimation with the novel matrix-F prior distribution (Mulder & Pericchi, 2018)
  - [Estimation](#) (Williams, 2018)
2. Bayesian hypothesis testing with the matrix-F prior distribution (Williams & Mulder, 2019)
  - [Exploratory hypothesis testing](#)
  - [Confirmatory hypothesis testing](#)
3. Comparing Gaussian graphical models (Williams, 2018; Williams, Rast, Pericchi, & Mulder, 2020)

- [Partial correlation differences](#)
- [Posterior predictive check](#)
- [Exploratory hypothesis testing](#)
- [Confirmatory hypothesis testing](#)

4. Extending inference beyond the conditional (in)dependence structure (Williams, 2018)

- [Predictability](#)(e.g., Haslbeck & Waldorp, 2018)
- [Posterior uncertainty intervals](#) for the partial correlations
- [Custom Network Statistics](#)

## Supported Data Types

- **Continuous:** The continuous method was described in Williams (2018). Note that this is based on the customary [Wishart distribution](#).
- **Binary:** The binary method builds directly upon Talhouk, Doucet, & Murphy (2012) that, in turn, built upon the approaches of Lawrence, Bingham, Liu, & Nair (2008) and Webb & Forster (2008) (to name a few).
- **Ordinal:** The ordinal methods require sampling thresholds. There are two approaches included in **BGGM**. The customary approach described in Albert & Chib (1993) (the default) and the ‘Cowles’ algorithm described in Cowles (1996).
- **Mixed:** The mixed data (a combination of discrete and continuous) method was introduced in Hoff (2007). This is a semi-parametric copula model (i.e., a copula GGM) based on the ranked likelihood. Note that this can be used for *only* ordinal data (not restricted to “mixed” data).

The computationally intensive tasks are written in c++ via the R package **Rcpp** (Eddelbuettel et al., 2011) and the c++ library **Armadillo** (Sanderson & Curtin, 2016). The Bayes factors are computed with the R package **BFpack** (Mulder et al., 2019). Furthermore, there are [plotting](#) functions for each method, control variables can be included in the model (e.g., `~ gender`), and there is support for missing values (see `bggm_missing`).

## Comparison to Other Software

**BGGM** is the only R package to implement all of these algorithms and methods. The mixed data approach is also implemented in the package **sbgcop** (base R, Hoff, 2007). The R package **BDgraph** implements a Gaussian copula graphical model in c++ (Mohammadi & Wit, 2015), but not the binary or ordinal approaches. Furthermore, **BGGM** is the only package for confirmatory testing and comparing graphical models with the methods described in Williams et al. (2020).

## Acknowledgements

DRW was supported by a National Science Foundation Graduate Research Fellowship under Grant No. 1650042 and JM was supported by a ERC Starting Grant (758791).

## References

- Albert, J. H., & Chib, S. (1993). Bayesian analysis of binary and polychotomous response data. *Journal of the American statistical Association*, 88(422), 669–679. doi:[10.1080/01621459.1993.10476321](https://doi.org/10.1080/01621459.1993.10476321)
- Chu, J.-h., Weiss, S. T., Carey, V. J., & Raby, B. A. (2009). A graphical model approach for inferring large-scale networks integrating gene expression and genetic polymorphism. *BMC systems biology*, 3(1), 55. doi:[10.1186/1752-0509-3-55](https://doi.org/10.1186/1752-0509-3-55)
- Cowles, M. K. (1996). Accelerating monte carlo markov chain convergence for cumulative-link generalized linear models. *Statistics and Computing*, 6(2), 101–111. doi:[10.1007/bf00162520](https://doi.org/10.1007/bf00162520)
- Eddelbuettel, D., François, R., Allaire, J., Ushey, K., Kou, Q., Russel, N., Chambers, J., et al. (2011). Rcpp: Seamless r and c++ integration. *Journal of Statistical Software*, 40(8), 1–18.
- Gelman, A., Meng, X.-L., & Stern, H. (1996). Posterior predictive assessment of model fitness via realized discrepancies. *Statistica Sinica*, 6(4), 733–807.
- Haslbeck, J. M., & Waldorp, L. J. (2018). How well do network models predict observations? On the importance of predictability in network models. *Behavior research methods*, 50(2), 853–861. doi:[10.3758/s13428-017-0910-x](https://doi.org/10.3758/s13428-017-0910-x)
- Hoff, P. D. (2007). Extending the rank likelihood for semiparametric copula estimation. *The Annals of Applied Statistics*, 1(1), 265–283. doi:[10.1214/07-AOAS107](https://doi.org/10.1214/07-AOAS107)
- Jeffreys, H. (1961). *The theory of probability*. Oxford: Oxford University Press. ISBN: [0191589675](https://www.isbn-international.org/en/number/0191589675)
- Kass, R. E., & Raftery, A. E. (1995). Bayes Factors. *Journal of the American Statistical Association*, 90(430), 773–795.
- Lawrence, E., Bingham, D., Liu, C., & Nair, V. N. (2008). Bayesian inference for multivariate ordinal data using parameter expansion. *Technometrics*, 50(2), 182–191. doi:[10.1198/004017008000000064](https://doi.org/10.1198/004017008000000064)
- Millington, T., & Niranjan, M. (2020). Partial correlation financial networks. *Applied Network Science*, 5(1), 11. doi:[10.1007/s41109-020-0251-z](https://doi.org/10.1007/s41109-020-0251-z)
- Mohammadi, R., & Wit, E. C. (2015). BDgraph: An r package for bayesian structure learning in graphical models. *Journal of Statistical Software*, 89(3). doi:[10.18637/jss.v089.i03](https://doi.org/10.18637/jss.v089.i03)
- Mulder, J., Gu, X., Olsson-Collentine, A., Tomarken, A., Böing-Messing, F., Hoijtink, H., Meijerink, M., et al. (2019). BFPack: Flexible bayes factor testing of scientific theories in r. *arXiv preprint arXiv:1911.07728*.
- Mulder, J., & Pericchi, L. (2018). The Matrix-F Prior for Estimating and Testing Covariance Matrices. *Bayesian Analysis*, (4), 1–22. doi:[10.1214/17-BA1092](https://doi.org/10.1214/17-BA1092)
- Rodriguez, J. E., Williams, D. R., Rast, P., & Mulder, J. (2020). On formalizing theoretical expectations: Bayesian testing of central structures in psychological networks. *PsyArXiv*. doi:[10.31234/osf.io/zw7pf](https://doi.org/10.31234/osf.io/zw7pf)
- Rubin, D. B. (1984). Bayesianly justifiable and relevant frequency calculations for the applied statistician. *The Annals of Statistics*, 1151–1172. doi:[10.1214/aos/1176346785](https://doi.org/10.1214/aos/1176346785)
- Sanderson, C., & Curtin, R. (2016). Armadillo: A template-based c++ library for linear algebra. *Journal of Open Source Software*, 1(2), 26. doi:[10.21105/joss.00026](https://doi.org/10.21105/joss.00026)
- Talhouk, A., Doucet, A., & Murphy, K. (2012). Efficient bayesian inference for multivariate probit models with sparse inverse correlation matrices. *Journal of Computational and Graphical Statistics*, 21(3), 739–757. doi:[10.1080/10618600.2012.679239](https://doi.org/10.1080/10618600.2012.679239)

- Webb, E. L., & Forster, J. J. (2008). Bayesian model determination for multivariate ordinal and binary data. *Computational statistics & data analysis*, 52(5), 2632–2649. doi:[10.1016/j.csda.2007.09.008](https://doi.org/10.1016/j.csda.2007.09.008)
- Williams, D. R. (2018). Bayesian Estimation for Gaussian Graphical Models: Structure Learning, Predictability, and Network Comparisons. *arXiv*. doi:[10.31234/OSF.IO/X8DPR](https://doi.org/10.31234/OSF.IO/X8DPR)
- Williams, D. R., & Mulder, J. (2019). Bayesian Hypothesis Testing for Gaussian Graphical Models: Conditional Independence and Order Constraints. *PsyArXiv*. doi:[10.31234/osf.io/ypxd8](https://doi.org/10.31234/osf.io/ypxd8)
- Williams, D. R., Rast, P., Pericchi, L. R., & Mulder, J. (2020). Comparing gaussian graphical models with the posterior predictive distribution and bayesian model selection. *Psychological Methods*. doi:[10.1037/met0000254](https://doi.org/10.1037/met0000254)
- Zerenner, T., Friederichs, P., Lehnertz, K., & Hense, A. (2014). A gaussian graphical model approach to climate networks. *Chaos: An Interdisciplinary Journal of Nonlinear Science*, 24(2), 023103. doi:[10.1063/1.4870402](https://doi.org/10.1063/1.4870402)